

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 9/28, 31/44, 9/20</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/09092</b>
			(43) International Publication Date: 24 February 2000 (24.02.00)
<p>(21) International Application Number: PCT/EP99/05724</p> <p>(22) International Filing Date: 7 August 1999 (07.08.99)</p> <p>(30) Priority Data: 98115141.8 12 August 1998 (12.08.98) EP</p> <p>(71) Applicant (for all designated States except US): BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH [DE/DE]; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): DIETRICH, Rango [DE/DE]; Im Tiergarten 16, D-78465 Konstanz (DE). NEY, Hartmut [DE/DE]; Peter-Thumb-Strasse 46, D-78464 Konstanz (DE).</p> <p>(74) Common Representative: BYK GULDEN LOMBERG CHEMISCHE FABRIK GMBH; Byk-Gulden-Strasse 2, D-78467 Konstanz (DE).</p>			<p>(81) Designated States: AE, AL, AU, BA, BG, BR, CA, CN, CZ, EE, GE, HR, HU, ID, IL, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, US, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> With international search report.</p>
<p>(54) Title: ORAL ADMINISTRATION FORM FOR PYRIDIN-2-YLMETHYLSULFINYL-1H-BENZIMIDAZOLES</p> <p>(57) Abstract</p> <p>The invention relates to an oral administration form for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles and their salts, which comprises the active compound together with tablet disintegrants and is provided with a film coating customary per se for sustained-release compositions.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ORAL ADMINISTRATION FORM FOR PYRIDIN-2-YLMETHYLSULFINYL-1H-BENZIMIDAZOLES

**Subject of the invention**

The present invention relates to a novel oral administration form for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles.

**Prior art**

Pyridin-2-ylmethylsulfinyl-1H-benzimidazoles and compounds structurally related to these, such as are disclosed, for example, in EP-A-0005129, EP-A-0166287, EP-A-0174726, EP-A-0268956, DE-A-3531487 and EP-A-0434999, have, on account of their H<sup>+</sup>/K<sup>+</sup>ATPase-inhibiting action, considerable importance in the therapy of diseases which are due to increased gastric acid secretion. Examples of active compounds from this group which are commercially available or in an advanced stage of clinical testing are 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulfinyl]-1H-benzimidazole (INN: omeprazole), 5-methoxy-2-[(S)-(4-methoxy-3,5-dimethyl-2-pyridinyl)methylsulfinyl]-1H-benzimidazole (prop. INN: esomeprazole), 5-difluoromethoxy-2-[(3,4-dimethoxy-2-pyridinyl)methylsulfinyl]-1H-benzimidazole (INN: pantoprazole), 2-[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridinyl)methyl-sulfinyl]-1H-benzimidazole (INN: lansoprazole), 2-[[4-(3-methoxypropoxy)-3-methylpyridin-2-yl]methyl-sulfinyl]-1H-benzimidazole (INN: rabeprazole), 2-[2-(N-isobutyl-N-methylamino)benzylsulfinyl]benzimidazole (lémiprazole) and 2-(4-methoxy-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridin-9-ylsulfinyl)-1H-benzimidazole (nepaprazole).

A common characteristic of the abovementioned pyridin-2-ylmethylsulfinyl-1H-benzimidazoles is the acid sensitivity - which is finally indispensable for their efficacy - of these active compounds, which is seen in their strong tendency to decompose in a neutral and, in particular, acidic environment, strongly colored decomposition products being formed.

In the past, there have been considerable efforts, despite the acid sensitivity of the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles, to obtain stable and storable oral administration forms which contain these compounds. There have likewise been efforts to obtain custom administration forms for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles for certain application purposes.

European Patent EP-B1-244 380 claims an oral administration form for certain pyridin-2-ylmethylsulfinyl-1H-benzimidazoles in which the active compound present in the tablet or pellet core is protected from the gastric acid by an enteric coating, a water-soluble intermediate layer which is intended to protect the core and acidic coating from one another additionally being situated between the active compound core and enteric coating.

- 2 -

The protection of the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles from gastric acid by application of an enteric coating can be regarded as the method of choice up to now when oral administration forms for this class of active compound are involved. The enteric coatings, whose resistance to gastric juice is based on the fact that free acidic groups (in particular carboxyl groups) are present in a polymer, must be separated, however, from the acid-sensitive active compound cores by suitable measures. This is carried out by application or production of a protective intermediate layer composed in whatever way (see, for example, EP-B1-589 981, WO-A-9601624, WO-A-9623500, WO-A-9624338, WO-A-9402140, WO-A-9712580 and WO-A-9800115).

### Description of the invention

Surprisingly, it has now been found that an enteric coating for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles is unnecessary if the coating used instead of it is designed so that the active compound is released only after a defined time, namely after gastric passage. Furthermore, it has surprisingly been found that, with a suitable design of the core comprising the active compound, the release of the active compound - once it has commenced - takes place within a short space of time, so that a rapidly rising and high active compound blood level is achieved.

The invention thus relates to an oral administration form for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles and their salts, which comprises the active compound together with tablet disintegrants and is provided with a film coating which is customary per se for sustained-release compositions.

Possible oral administration forms are, for example, pellets, microtablets, minitablets or in particular tablets, if desired dispensed in capsules.

Suitable pyridin-2-ylmethylsulfinyl-1H-benzimidazoles within the meaning of the invention are, for example, omeprazole, esomeprazole, lansoprazole, rabeprazole, leminoprazole, nepaprazole and in particular pantoprazole.

Salts of the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles which may be mentioned primarily are the salts with bases, in particular the sodium, potassium, calcium and magnesium salt. The pantoprazole sodium salts, in particular the pantoprazole sodium sesquihydrate, is particularly preferred.

Possible tablet disintegrants are the customary agents known to the person skilled in the art. Examples which may be mentioned are certain cellulose derivatives (e.g. sodium cellulose glycolate and Tyloses), starch, compositions based on sodium carboxymethylcellulose and potato starch (e.g. Primojel), sodium carboxymethylstarch (e.g. Explotab), bentonite, sodium alginate or pectin, but in particular chemically indifferent agents such as crosslinked polyvinylpyrrolidone (e.g. Crospovidone). The content of tablet disintegrant is customarily between 2 and 10 % by weight based on the entire core. Depending

- 3 -

on the type of tablet disintegrant, however, larger contents can also be used, in the case of Crospovidone, for example, 20-35% by weight.

In addition to the tablet disintegrant, if desired the tablet cores contain further auxiliaries and fillers or binders. Auxiliaries used are, in particular, lubricants and release agents. Mention may be made here, for example, of calcium salts of higher fatty acids, such as, for example, calcium stearate. Binders which may be mentioned are, in particular, polyvinylpyrrolidone and/or hydroxypropylmethylcellulose and, if desired, mannitol, which is additionally preferred as a filler.

To increase the stability of the tablet cores, it has proven advantageous to employ the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles in the form of their salts and/or with addition of one or more physiologically tolerable inorganic compounds having a basic reaction. Mention may be made here, for example, of the pharmacologically tolerable alkali metal, alkaline earth metal or earth metal salts of weak acids and the pharmacologically tolerable hydroxides and oxides of alkaline earth metals and earth metals. A base to be emphasized by way of example which may be mentioned is sodium carbonate.

Film coatings customary for sustained-release compositions which may be mentioned are membranes made of plastics having a low swelling power in water, in which small soluble particles are embedded, or in particular those swellable plastic membranes which contain a small proportion of a suitable salt which determines the permeability of the film coating.

Plastics suitable for the construction of the membranes are those which are water-insoluble and physiologically tolerable. Plastics having a low swelling power in water are understood for the purposes of the present invention as meaning, for example, those which absorb not more than 5% by weight of water in aqueous medium. For this, cellulose ethers and cellulose esters are regarded as particularly suitable. In addition, suitable plastics are also polymers such as polyvinyl chloride. Swellable plastics which may be mentioned are, in particular, copolymers of acrylic and methacrylic acid esters.

Small soluble particles which may be mentioned are, for example, lactose crystals, which are preferably employed in micronized form. The particle size is expediently less than 20  $\mu\text{m}$ , preferably less than 10  $\mu\text{m}$ . The ratio of plastic to soluble particles can be varied within wide limits. A weight ratio of plastic to soluble particles of approximately 2:1 to 1:3 is preferred. A weight ratio of 4:3 to 4:5 is particularly preferred.

Salts suitable for the swellable plastic membranes which may be mentioned are, for example, ammonium salts, in particular quaternary ammonium salts. In a particular embodiment of plastic membranes, some of the ester groups of a copolymer of acrylic and methacrylic acid esters are ester groups having quaternary ammonium structures. An example of such copolymers having quaternary ammonium

- 4 -

groups which may be mentioned is trimethylammonium methyl methacrylate chloride (e.g. Eudragit RL or Eudragit RS from Röhm).

The release time of the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles can be controlled within a wide range by variation of the composition of the membrane and/or by variation of the layer thickness of the membrane. Thus, release is effected at an earlier time by lowering the layer thickness of the membrane, by increasing the proportion of soluble particles, by use of the soluble particles in a more coarse-grained form or, in the case of the swellable plastic membranes, by increasing the proportion of a suitable salt (e.g. higher proportion of quaternary ammonium groups in the copolymer of acrylic and methacrylic acid esters).

The application of the membrane to the tablet cores is carried out in a manner known per se, in particular by one of the customary spraying techniques. For this, a solution of the plastic or plastic mixture intended for the membrane is prepared in a solvent or in a solvent mixture or preferably an aqueous dispersion of the plastic or plastic mixture. The soluble, micronized particles are suspended in the solution before the spraying. If necessary, the suspension is stirred during the spraying in order to prevent settling of the suspended particles. In the case of the preferred procedure using aqueous dispersions, the salts responsible for the permeability of the plastic are already contained in the plastic itself in the form of quaternary ammonium groups. In the case of application of the membrane from an aqueous dispersion, it is also possible to work under alkaline conditions.

The membrane can contain the customary auxiliaries, such as plasticizers, wetting agents, colorants and antiadherents. Pharmacologically tolerable plasticizers such as, for example, polyethylene glycols, paraffins, glycerol or propylene glycol are suitable. Wetting agents may be necessary if the coating is to be dyed with dye lakes. Sorbitol fatty acid esters or salts of dioctylsulfosuccinic acid, for example, are suitable. Antiadherents which may be mentioned are, in particular, calcium stearate or talc.

With respect to the preparation and construction of the tablet cores reference is made, for example, to the embodiments in European Patent EP-B1-589 981.

The following examples of administration forms according to the invention explain the invention in greater detail without restricting it.

**Examples****Example 1: Tablets****A. Tablet cores with 10 mg of active compound**

	Ingredients	per core
(a)	pantoprazole Na $\times$ 1.5 H <sub>2</sub> O	11.28 mg
(b)	sodium carbonate, anhydrous	2.50 mg
(c)	mannitol	10.68 mg
(d)	PVP, insoluble (Crospovidone)	12.50 mg
(e)	PVP 90	1.00 mg
(f)	calcium stearate	0.80 mg
	Total per core	38.75 mg

(a) is mixed with some of (b), (c) and (d). The remainder of (b) and (c) is added to the clear aqueous solution of (e) and adjusted to a pH of  $> 10$  using (b). Granulation is carried out in a fluidized bed granulator using this solution. The remainder of (d) and (f) is added to the dried granules and the granules are pressed in a suitable tablet machine.

**B. Coating**

	Ingredients	Initial weight	Coating per core
(g)	Eudragit RS 30 D	2400.00 g	4.876 mg
(h)	purified water	4800.00 g	
(i)	propylene glycol	144.00 g	0.975 mg
(j)	Ca stearate	21.60 g	0.146 mg
(k)	1 N NaOH	81.10 g	0.002 mg
	Total film coating	7446.70 g	6.000 mg

The ingredients are stirred to give a dispersion which is screened before processing. The dispersion is sprayed onto the cores obtained under A in a suitable apparatus.

The coating application of 6 mg per tablet core leads to a spontaneously commencing and complete release of active compound after 2 hours.

**Example 2: Combinations**

The following combinations of tablets according to the invention (prepared according to Example 1, comprising 10 mg of active compound, below "tablet E") and the known enteric tablets (prepared according to EP-B-589981, comprising 10 mg of active compound, below "tablet M") are, for example, conceivable, the tablets being dispensed into hard gelatin capsules of size 3:

- 1 tablet E + 1 tablet M
- 2 tablets E + 2 tablets M
- 3 tablets E + 1 tablet M
- 1 tablet E + 3 tablets M

Instead of the enteric tablets, the pellets prepared according to EP-B-589981 can also be used.



**Commercial applicability**

The oral administration forms according to the invention can be employed for the treatment and prevention of all the diseases which are considered to be treatable or avoidable by the use of pyridin-2-ylmethylsulfinyl-1H-benzimidazoles. In particular, the oral administration forms according to the invention can be employed in the treatment of disorders of the stomach.

Surprisingly, sustained (i.e. more or less constant over a relatively long period) release behavior is not achieved using the oral administration forms according to the invention - despite the use of a customary sustained-release coating. On the contrary, initially no active compound at all is released over a certain period, the length of this period - as explained above - being controllable by the type and thickness of the membrane.

After expiry of the adjustable period, all of the active compound is then released within a very short space of time. Due to the dissolution of the particles embedded in the membrane, the membrane becomes porous or, due to the swelling of the permeable membrane, this becomes permeable and water penetrates into the core; as a result of this the tablet disintegrant begins to swell, and when the swelling pressure is sufficient in order to disintegrate the membrane, the active compound is released spontaneously and completely.

With the aid of the oral administration form according to the invention, it is thus possible to simulate an administration of active compound at a later time. As a result, the possibility is opened up of allowing a once daily administration instead of a twice daily administration of the active compound to begin by combining, for example, in one and the same administration form (e.g. in a capsule) two active compound forms whose release is different (e.g. a customary, enteric tablet and a tablet according to the invention).

The invention therefore further relates to the combination of an oral administration form according to the invention with a conventional (i.e. enteric-coated) administration form for pyridin-2-ylmethylsulfinyl-1H-benzimidazoles. "Combination" in this connection is understood as meaning the fixed or free combination.

In the fixed combination, both administration forms are present in a single dose unit (e.g. in a common tablet of outer conventional construction and inner core coated according to the invention, in a capsule comprising conventionally coated pellets and pellets according to the invention, or in particular in a capsule comprising two or more tablets, of which at least one corresponds to the specification according to the invention).

- 8 -

In the free combination, the two administration forms (that according to the invention and the conventional one) are present in separate dose units, which can be contained in a common packaging unit or in separate packaging units. In a common packaging unit, the different administration forms, for example, can be arranged in the form of capsules or tablets in rows lying next to one another in a blister pack. At the time indicated by the physician, the patient would in each case successively take a capsule or tablet from each row within a short length of time (in particular within 5 minutes).

Independently of whether a fixed or free combination is present, the compliance in the case of the combination according to the invention is in any case considerably greater than when two conventional administration forms have to be taken in a relatively large space of time (for example in the space of 3 to 12 hours).

The two-fold administration of active compound simulated by the fixed or free combination leads in a relatively large space of time (compared with the same dose of active compound as a single administration) to a smaller width of variation in the active compound blood levels in the patients and moreover to more rapid symptom relief.

In this connection, the fixed combination is preferred, particularly the combination of pellets according to the invention and conventional pellets and very particularly the combination of tablets according to the invention and conventional tablets in one capsule.

The treatment dose for an adult patient is, with respect to the pyridin-2-ylmethylsulfinyl-1H-benzimidazoles or their pharmaceutically tolerable salts, approximately 5 mg to 100 mg, in particular 10 mg to 80 mg, preferably 20 mg to 40 mg per day, calculated on the free acid. This treatment dose can be evenly or unevenly divided over the two administration forms in the combination according to the invention. A more or less equal division is preferred, e.g. 20 mg of the administration form according to the invention and 20 mg of the conventional (enteric-coated) administration form, in each case based on the free acid.

For their part, the oral administration forms according to the invention or the combinations according to the invention can in turn be combined with other medicaments, in particular with antimicrobial agents, such as are employed for the control of the bacterium *Helicobacter pylori* (*H. pylori*). Suitable antimicrobial agents for the control of the bacterium *H. pylori* which may be mentioned are bismuth salts [e.g. bismuth subcitrate, bismuth subsalicylate, ammonium bismuth(III) potassium citrate dihydroxide, bismuth nitrate oxide, dibismuth tris(tetraoxodialuminate)], but in particular  $\beta$ -lactam antibiotics, for example penicillins (such as benzylpenicillin, phenoxymethylpenicillin, propicillin, azidocillin, dicloxacillin, flucloxacillin, oxacillin, amoxycillin, bacampicillin, ampicillin, mezlocillin, piperacillin or azlocillin), cephalosporins (such as cefadroxil, cefaclor, cefalexin, cefixime, cefuroxime, cefatamet, cefadroxil, ceftibuten, cefpodoxime, cefotetan, cefazolin, cefoperazone, ceftizoxime, cefotaxime, ceftazidime,

- 9 -

cefamandol, cefepime, cefoxitin, cefodizime, cefsulodin, ceftriaxone, cefotiam or cefmenoxime) or other  $\beta$ -lactam antibiotics (e.g. aztreonam, loracarbef or meropenem); enzyme inhibitors, for example sulbactam; tetracyclines, for example tetracycline, oxytetracycline, minocycline or doxycycline; aminoglycosides, for example tobramycin, gentamicin, neomycin, streptomycin, amikacin, netilmicin, paromomycin or spectinomycin; amphenicols, for example chloramphenicol or thiamphenicol; lincomycins and macrolide antibiotics, for example clindamycin, lincomycin, erythromycin, clarithromycin, spiramycin, roxithromycin or azithromycin; polypeptide antibiotics, for example colistin, polymixin B, teicoplanin or vancomycin; gyrase inhibitors, for example norfloxacin, cinoxacin, ciprofloxacin, pipemidic acid, enoxacin, nalidixic acid, pefloxacin, fleroxacin or ofloxacin; nitroimidazoles, for example metronidazole; or other antibiotics, for example fosfomycin or fusidic acid, where these antibacterially active substances - together with the oral administration forms according to the invention or with the combinations according to the invention - can be administered on their own or alternatively combined with one another. Combinations of antibacterially active substances which may be mentioned are, for example, amoxicillin plus metronidazole, clarithromycin plus metronidazole and amoxicillin plus clarithromycin.

**Patent claims**

1. An oral administration form for pyridin-2-ylmethylsulfinyl-1H-benzimidazole and its salts, which comprises the active compound together with tablet disintegrants and is provided with a film coating-customary per se for sustained which is release compositions.
2. The administration form as claimed in claim 1, wherein the pyridin-2-ylmethylsulfinyl-1H-benzimidazole is omeprazole, esomeprazole, lansoprazole, rabeprazole, leminoprazole or nepaprazole.
3. The administration form as claimed in claim 1, wherein the pyridin-2-ylmethylsulfinyl-1H-benzimidazole is pantoprazole.
4. The administration form as claimed in claim 1, wherein the tablet disintegrant is Crospovidone.
5. The administration form as claimed in claim 1, wherein the tablet disintegrant is Crospovidone having a proportion in the tablet core of 20-35% by weight.
6. The administration form as claimed in claim 1, wherein the film coating is a copolymer of acrylic and methacrylic acid esters having quaternary ammonium structures.
7. A combination consisting of an administration form as claimed in claim 1 and an administration form of a pyridin-2-ylmethylsulfinyl-1H-benzimidazole having an enteric coating.
8. The administration form as claimed in claim 1 in combination with or for combined use with an antimicrobial agent.
9. The combination as claimed in claim 7 in combination with or for combined use with an antimicrobial agent.
10. The use of administration forms and combinations as claimed in one of claims 1 to 9 in the treatment of disorders of the stomach.

# INTERNATIONAL SEARCH REPORT

International Application No

PC./EP 99/05724

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/28 A61K31/44 A61K9/20

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, Y, L	WO 97 02020 A (BYK GULDEN LOMBERG CHEM FAB) 23 January 1997 (1997-01-23) the whole document ---	1-10
Y	WO 97 25979 A (PERIO PROD LTD) 24 July 1997 (1997-07-24) the whole document ---	1, 6, 10
Y	EP 0 519 365 A (BYK GULDEN LOMBERG CHEM FAB) 23 December 1992 (1992-12-23) cited in the application the whole document ---	1-10
Y	EP 0 793 959 A (TAKEDA CHEMICAL INDUSTRIES LTD) 10 September 1997 (1997-09-10) the whole document ---	1-10
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

26 October 1999

Date of mailing of the international search report

05/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fischer, W

# INTERNATIONAL SEARCH REPORT

International Application No

PC./EP 99/05724

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 42 19 390 A (BYK GULDEN LOMBERG CHEM FAB) 24 December 1992 (1992-12-24) the whole document ---	1-5, 10
A	EP 0 526 862 A (VECTORPHARMA INT) 10 February 1993 (1993-02-10) -----	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05724

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9702020	A	23-01-1997	US 5945124 A AU 6517496 A CA 2232450 A EP 0841903 A JP 11508577 T	31-08-1999 05-02-1997 23-01-1997 20-05-1998 27-07-1999
WO 9725979	A	24-07-1997	US 5840332 A AU 1206597 A CN 1208343 A CZ 9802198 A EP 0877604 A	24-11-1998 11-08-1997 17-02-1999 16-12-1998 18-11-1998
EP 0519365	A	23-12-1992	AT 144416 T AU 683411 B AU 1974692 A BG 61796 B BG 98286 A CA 2109697 A CN 1067809 A,B CZ 9302764 A DE 4219390 A DE 59207438 D DK 589981 T WO 9222284 A EP 0589981 A ES 2096080 T FI 935677 A GR 3022154 T HK 1005851 A HR 920162 A IE 77640 B IL 102096 A JP 6508118 T LV 11982 A LV 11982 B MX 9202961 A NO 934648 A NZ 243147 A PL 169951 B RU 2089180 C SK 128793 A ZW 9392 A	15-11-1996 13-11-1997 12-01-1993 30-06-1998 15-08-1994 23-12-1992 13-01-1993 13-07-1994 24-12-1992 28-11-1996 17-03-1997 23-12-1992 06-04-1994 01-03-1997 16-12-1993 31-03-1997 29-01-1999 31-08-1996 31-12-1997 18-06-1996 14-09-1994 20-03-1998 20-09-1998 01-02-1993 16-12-1993 21-12-1995 30-09-1996 10-09-1997 08-06-1994 17-02-1993
EP 0793959	A	10-09-1997	CA 2199345 A CN 1164424 A JP 9295933 A	07-09-1997 12-11-1997 18-11-1997
DE 4219390	A	24-12-1992	AT 144416 T AU 683411 B AU 1974692 A BG 61796 B BG 98286 A CA 2109697 A CN 1067809 A,B CZ 9302764 A DE 59207438 D DK 589981 T WO 9222284 A	15-11-1996 13-11-1997 12-01-1993 30-06-1998 15-08-1994 23-12-1992 13-01-1993 13-07-1994 28-11-1996 17-03-1997 23-12-1992

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05724

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 4219390 A		EP 0519365 A	23-12-1992
		EP 0589981 A	06-04-1994
		ES 2096080 T	01-03-1997
		FI 935677 A	16-12-1993
		GR 3022154 T	31-03-1997
		HK 1005851 A	29-01-1999
		HR 920162 A	31-08-1996
		IE 77640 B	31-12-1997
		IL 102096 A	18-06-1996
		JP 6508118 T	14-09-1994
		LV 11982 A	20-03-1998
		LV 11982 B	20-09-1998
		MX 9202961 A	01-02-1993
		NO 934648 A	16-12-1993
		NZ 243147 A	21-12-1995
		PL 169951 B	30-09-1996
		RU 2089180 C	10-09-1997
		SK 128793 A	08-06-1994
		ZW 9392 A	17-02-1993
EP 0526862 A	10-02-1993	IT 1251153 B	04-05-1995
		AT 134134 T	15-02-1996
		DE 69208299 D	28-03-1996
		DE 69208299 T	18-07-1996
		ES 2086029 T	16-06-1996